

Chapter 10 FRQ Homework

1.

For $t \geq 0$, a particle is moving along a curve so that its position at time t is $(x(t), y(t))$. At time $t = 2$, the particle is at position $(1, 5)$. It is known that $\frac{dx}{dt} = \frac{\sqrt{t+2}}{e^t}$ and $\frac{dy}{dt} = \sin^2 t$.

- Is the horizontal movement of the particle to the left or to the right at time $t = 2$? Explain your answer. Find the slope of the path of the particle at time $t = 2$.
- Find the x -coordinate of the particle's position at time $t = 4$.
- Find the speed of the particle at time $t = 4$. Find the acceleration vector of the particle at time $t = 4$.
- Find the distance traveled by the particle from time $t = 2$ to $t = 4$.

2.

The polar curve r is given by $r(\theta) = 3\theta + \sin \theta$, where $0 \leq \theta \leq 2\pi$.

- Find the area in the second quadrant enclosed by the coordinate axes and the graph of r .
- For $\frac{\pi}{2} \leq \theta \leq \pi$, there is one point P on the polar curve r with x -coordinate -3 . Find the angle θ that corresponds to point P . Find the y -coordinate of point P . Show the work that leads to your answers.
- A particle is traveling along the polar curve r so that its position at time t is $(x(t), y(t))$ and such that $\frac{d\theta}{dt} = 2$. Find $\frac{dy}{dt}$ at the instant that $\theta = \frac{2\pi}{3}$, and interpret the meaning of your answer in the context of the problem.

3.

The velocity vector of a particle moving in the plane has components given by

$$\frac{dx}{dt} = 14 \cos(t^2) \sin(e^t) \quad \text{and} \quad \frac{dy}{dt} = 1 + 2 \sin(t^2), \quad \text{for } 0 \leq t \leq 1.5.$$


At time $t = 0$, the position of the particle is $(-2, 3)$.

- For $0 < t < 1.5$, find all values of t at which the line tangent to the path of the particle is vertical.
- Write an equation for the line tangent to the path of the particle at $t = 1$.
- Find the speed of the particle at $t = 1$.
- Find the acceleration vector of the particle at $t = 1$.

4.

The graphs of the polar curves $r = 2$ and $r = 3 + 2 \cos \theta$ are shown in the figure above. The curves intersect when $\theta = \frac{2\pi}{3}$ and $\theta = \frac{4\pi}{3}$.

- Let R be the region that is inside the graph of $r = 2$ and also inside the graph of $r = 3 + 2 \cos \theta$, as shaded in the figure above. Find the area of R .
- A particle moving with nonzero velocity along the polar curve given by $r = 3 + 2 \cos \theta$ has position $(x(t), y(t))$ at time t , with $\theta = 0$ when $t = 0$. This particle moves along the curve so that $\frac{dr}{dt} = \frac{dr}{d\theta}$.



Find the value of $\frac{dr}{dt}$ at $\theta = \frac{\pi}{3}$ and interpret your answer in terms of the motion of the particle.

- For the particle described in part (b), $\frac{dy}{dt} = \frac{dy}{d\theta}$. Find the value of $\frac{dy}{dt}$ at $\theta = \frac{\pi}{3}$ and interpret your answer in terms of the motion of the particle.